

Table 1. Baseline characteristics of the patients	
After TAVR PPM implant (n=17)	
Age (years)	74.7 ± 9
Women (%)	10 (58.8%)
Diabetes (%)	6 (35.2%)
Hypertension (%)	13 (76.4%)
LVEF (%)	56.9 ± 15.3
EF <35 (%)	1 (5.8%)
EF 35-50 (%)	4 (23.5%)
EF >50 (%)	12 (70.5%)
Functional class (NYHA)	
1	1 (0%)
2	11 (64.7%)
3	6 (35.2%)
4	0 (0%)
Coronary artery disease	6 (35.2%)
PR (ms)	176.4 ± 27.6
PR >200ms	6 (35.2%)
QRS (ms)	86.4 ± 15.7
QRS >120ms	2 (11.7%)
DM2: diabetes, LVEF: left ventricular ejection fraction, PR: PR interval, QRS: QRS complex, ms: millisecond.	

Table 2. Prosthesis type and PPM indication	
After TAVR PPM implant (n=17)	
LVOT depth (mm)	5.5 ± 2.9
LBBB (%)	7 (41.1%)
HAVB (%)	5 (29.4%)
LBBB + HAVB (%)	5 (29.4%)
Corevalve (%)	6 (35.2%)
Size (mm)	29 ± 2.4
LVOT depth (mm)	7 ± 3.9
BRIHH (%)	2 (33.3%)
HAVB (%)	2 (33.3%)
LBBB + HAVB (%)	2 (33.3%)
Lotus (%)	4 (23.5%)
Size (mm)	25 ± 1.6
LVOT depth (mm)	4 ± 0.8
LBBB (%)	2 (50%)
HAVB (%)	1 (25%)
LBBB + HAVB (%)	1 (25%)
Evolut R (%)	3 (17.6%)
Size (mm)	28.6 ± 4.6
LVOT depth (mm)	5.3 ± 4
LBBB (%)	1 (33.3%)
HAVB (%)	1 (33.3%)
LBBB + HAVB (%)	1 (33.3%)
Pórtico (%)	3 (17.6%)
Size (mm)	25 ± 1

LVOT depth (mm)	5 ± 1
LBBB (%)	1 (33.3%)
HAVB (%)	1 (33.3%)
LBBB + HAVB (%)	1 (33.3%)
Edwards (%)	1 (5.8%)
PPM: permanent pacemaker, LVOT: left ventricular outflow tract, LBBB: left bundle-branch block, HAVB: high-degree atrioventricular block.	

Table 3. Cardiac stimulation percentage	
After TAVR PPM implant (n=17)	
Global auricular stimulation (%)	13.4 ± 16.6
Global ventricular stimulation (%)	71.7 ± 42.6
Ventricular stimulation	
LBBB indication (%)	72.4
HAVB indication (%)	99.2 ± 0.5
LBBB + HAVB indication (%)	52.5
Progression from LBBB to HAVB (%)	3 (17.6%)
Recovery after HAVB (%)	3 (17.6%)
Ventricular stimulation	
Corevalve (%)	85
Lotus (%)	99 ± 0.8
Portico (%)	66 ± 0
Evolute R (%)	38.3
LBBB: left bundle-branch block, HAVB: high-degree atrioventricular block.	